



**AMINOCAL**  
*improving the firmness  
of green tissues and fruits*

**Aminocal** is formulated with enzymatically hydrolyzed, biologically active free **Amino acids** and a **complex of Metabolites**, combined with naturally chelated **Calcium** to deliver yields that are higher in quality and quantity.

The natural Amino acids and Metabolites in **Aminocal** supports plant maintenance, growth, vitality and reproduction by supplying additional energy through the Krebs cycle and enhancing stress resistance.

The natural chelated Calcium corrects the main physiological disorders of calcium deficiency and improves the firmness of green tissues and fruits, extending the crop's shelf life. This results in improved plant and fruit colour, larger leaves, longer roots, more flowers, more fruit and/or larger fruit.

## Benefits

- **Triple action: Bio-stimulant, Fertilizer and Calcium deficiency corrector**
- **Fast and efficient leaf penetration of calcium**
- **Rapid recovery from critical situations**
- **Improves fruit set, quality and shelf life**
- **Reduces Calcium deficiency problems like bitterpit & blossom end rot**

Composition	(w/w)	(w/w)
<b>Free amino acids</b>	10%	Total amino acids
Organic matter	20%	Dry matter
Organic nitrogen	2.9%	Total nitrogen
Calcium (CaO)	5%	Ca
<b>Biologically Active Metabolites</b> (incl. Bile acids, Lipids, Carbohydrates, Nucleotides Peptides)		>150
Specific weight	1.075kg/liter	pH
		5-8

### Application rate

**Foliar application:** 500ml-1 l/ha (300-400mL/hL water)

- Apply in the early morning or late evening
- Clean equipment after use.

**Drip irrigation:** 2-3 l/ha

- Apply the first drip irrigation moments early in the morning so that the pipes are clean during the night.
- Repeat the application every 10-15 days during the growth season

### Application time

It is advisable to use it during Calcium deficiencies and when the crop needs a physiological stimulus, such as:

- Start of the vegetative cycle, as soon as it has sufficient leaf mass
- Pre-flowering and petal-falling phase
- Fruit development

**Aminocal** also prevents the risk of Bitter pit and Fruit cracking in pome fruits, blossom-end rot in tomatoes and solonaceous crops like tomatoes and tobacco, tip burn in leafy vegetables and leaf and stem spot in melons.

## Free amino acid composition g/kg product

Alanine	9.7
Arginine	2.8
Aspartic acid	9.1
Cystine	0.1
Glutamic acid	11.1
Glycine	6.5
Histidine	3.9
Hydroxyproline	0.1
Isoleucine	5.7
Leucine	10.3
Lysine	12.0
Methionine	3.0
Phenylalanine	7.7
Proline	7.7
Serine	6.9
Taurine	0.8
Threonine	7.7
Thryptophan	1.5
Tyrosine	1.3
Valine	8.8



## Packaging

IBC:	1000 L
BARREL:	220 L
CANNISTER:	5, 10 & 20 L
BOTTLES:	1 L

\*above mentioned figures are indicative

## Storage information

Product is stable when stored under normal undiluted conditions. The original product in the state in which it is delivered is hazard-free concerning the transmission of pathogens. Close package properly after the use, store in a frost-free place and avoid storage at direct sunlight! Shake well before use.

The product is produced in conformity with EU Directives 837/2007 and 889/2008 and is listed on the FiBL Betriebsmittelliste as approved for organic farming.

